

RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL AND DERMAL TOXICITY; AND BIRD TOXICITY. FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION.

SECTION 24(C) REGISTRATION

FOR DISTRIBUTION AND USE ONLY WITHIN MONTANA



PHORATE 20 G

**EPA REG. NO. 34704-259
EPA SLN NO. MT-910004**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS
ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**THIS LABEL MUST BE IN POSSESSION
OF USER AT THE TIME OF APPLICATION.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

POTATOES — For control of aphids, leafhoppers, leaf miners, psyllids, suppression of wireworms, flea beetle larvae, and reduction of flea beetle adults. On sandy or light soils, use 11¼ ozs. PHORATE 20 G per 1000 feet of row (minimum 32 inch spacing). On heavy or clay soils, use 17¼ ozs. PHORATE 20 G per 1000 feet of row (any row spacing). Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time, or at postemergence. Do not exceed 3 lbs. of the active ingredient Phorate per acre per year. Wait 90 days after treatment before harvesting potatoes. Do not apply within 200 feet of fish-bearing waters.

**24(C) Registrant:
PLATTE CHEMICAL CO.
419 18TH STREET
GREELEY, COLORADO 80631-5852**